

EUCLIDEAN COMPONENTS FOR A CLASS OF SELF-INJECTIVE ALGEBRAS

SARAH SCHEROTZKE (OXFORD)

ABSTRACT. We determine the length of composition series of projective modules of G -transitive algebras with an Auslander-Reiten component of Euclidean tree class. We thereby correct and generalize a result in [F2, 4.6]. Furthermore we show that modules with certain length of composition series are periodic. We apply these results to G -transitive blocks of the universal enveloping algebras of restricted p -Lie algebras and prove that G -transitive principal blocks only allow components with Euclidean tree class if $p = 2$. Finally we deduce conditions for a smash product of a local basic algebra Γ with a commutative semi-simple group algebra to have components with Euclidean tree class, depending on the components of the Auslander-Reiten quiver of Γ .

1. INTRODUCTION

The stable Auslander-Reiten quiver of a finite-dimensional algebra can be viewed as part of a presentation of the stable module category, and it is an important invariant which has many applications. It is a locally finite graph, where the vertices correspond to the isomorphism classes of indecomposable modules. Each connected component is isomorphic to $\mathbb{Z}[T]/\Gamma$ where T is a tree, and Γ is an admissible group of automorphism. (See [Ben1][4.15.6] for details).

For many self-injective algebras it is known that the possibilities for T are restricted, it can only be Dynkin, or Euclidean, or one of a few infinite trees (see [W], [E], [ES]). In this paper we study algebras with Euclidean components. Recently the study of self-injective algebras with Euclidean Auslander-Reiten components has attracted much attention, for example all self-injective algebras of Euclidean type have this property (see the survey article [Sk, Section 4]).

These have been also studied in the context of reduced enveloping algebras by Farnsteiner, and we discovered that [F2, 4.6] is not correct. In Theorem [F2, 4.6] Farnsteiner proves a necessary conditions for certain blocks of universal enveloping algebras $u(L, \chi)$ of a restricted p -Lie algebra L with $p > 2$, to have an Auslander-Reiten component with Euclidean tree class. Unfortunately one crucial step in the proof is wrong. As all other results in the case of $\tilde{D}_n$ -tree class depend on this step, we need a different proof and Theorem.

2000 *Mathematics Subject Classification.* 16G70.

Key words and phrases. Auslander-Reiten theory, G -transitive algebras .

In the first section we give a proof for the more general setup of G -transitive blocks of Frobenius algebras in any characteristic.

In the second section we apply the main result of the first section to G -transitive blocks of $u(L, \chi)$. We show that some of the results of Farnsteiner's paper remain true while others need additional assumptions. We can show that a G -transitive principal block of $u(L, \chi)$ does not have Auslander-Reiten components of Euclidean tree class if $p > 2$.

In the last section we determine conditions for the smash product of a basic local algebra Γ and a semi-simple commutative group algebra to have Auslander-Reiten components of Euclidean tree class depending on the tree class of components of Γ .

Let B be an indecomposable Frobenius algebra. We introduce for a group $G \subset \text{Aut}(B)$ the G -transitive algebra B . This means that G acts transitively on the set of simple modules in B . We denote by $l(P)$ the length of an indecomposable projective module P . We show in 3.3, 3.11, 3.7 that the following holds for G -transitive algebras that have Auslander-Reiten components of Euclidean tree class:

- (1) All non-periodic Auslander-Reiten components are either isomorphic to $\mathbb{Z}[\tilde{A}_{1,2}]$ or $\mathbb{Z}[\tilde{D}_n]$ for n odd and $n > 5$.
- (2) In the first case $l(P) = 4$ and all indecomposable modules of length $0 \bmod l(P)$ and $2 \bmod l(P)$ are periodic.
- (3) In the second case $l(P) = 8$ and all indecomposable modules of length $4 \bmod l(P)$ are periodic.

In [F2] Farnsteiner introduces $G(L)$, the group of group-like elements of $u^*(L)$ and shows that they can be embedded into the automorphism group of $u(L, \chi)$. He proves for $G(L)$ -transitive blocks over a field of characteristic $p > 2$ (see [F2, 4.6]) that:

- (1) All non-periodic Auslander-Reiten components are either isomorphic to $\mathbb{Z}[\tilde{A}_{1,2}]$ or $\mathbb{Z}[\tilde{D}_5]$. All indecomposable modules of length 2 are periodic.
- (2) In the first case $l(P) = 4$.
- (3) In the second case $l(P) = 8$.

For his results in the case of tree class $\tilde{D}_n$ he first shows that $n = 5$. As this step is wrong, a different proof was needed for the more general setup. In the new proof we can show that $n > 5$ which contradicts Farnsteiner's result.

Additionally we show that the number of non-periodic components is equal to the number of isomorphism types of simple modules in B .

In the second section we can verify Farnsteiner's statement that supersolvable algebras in characteristic $p > 2$ do not have Auslander-Reiten components of Euclidean tree class. An example by [E, 2.3] shows that this statement is wrong for $p = 2$. We also show in 4.2 that if B is a G -transitive

block of $u(L, \chi)$ that has an Auslander-Reiten component of Euclidean tree class, then $p = 2$ or the dimension of a simple module in B is divisible by 2.

In the last section we introduce the smash product of a basic local algebra Γ and a semi-simple group algebra kG , where $G \subset \text{Aut}(\Gamma)$ is an abelian group. We show that they are a special cases of G -transitive algebras and describe how to construct the Gabriel quiver of the smash product from the Gabriel quiver of Γ .

For a finite-dimensional algebra A , we introduce $G(A)$ the free abelian group with basis the isomorphism types of indecomposable modules. We have a bilinear form $(-, -)_A$ induced by $\dim_k \text{Hom}_A(-, -)$ defined on $G(A)$. We develop some properties of this bilinear form in 5.6, 5.7.

We use these general results to show in 5.15 that Auslander-Reiten sequences of the smash product restricted to Γ are sums of Auslander-Reiten sequences in Γ that are twists of each other. Using this result, we prove that the smash product has an Auslander-Reiten component of tree class $\tilde{D}_n$ if and only if Γ has an Auslander-Reiten component of tree class $\tilde{D}_n$. If the smash product has an Auslander-Reiten component of tree class $\tilde{A}_{1,2}$, then so does Γ . The converse is not true, as we show by providing a counter example. We can in this case restrict the component to $Z[\tilde{A}_{12}]$ and $\mathbb{Z}[\tilde{A}_n]$ for certain $n \in \mathbb{N}$ (see 5.17).

I would like to thank my Supervisor Karin Erdmann for her comments which have improved this a paper considerably. I would also like to thank the referee and the editor Andrzej Skowroński for their useful suggestions.

2. EUCLIDEAN COMPONENTS FOR G -TRANSITIVE ALGEBRAS

For general background on Auslander-Reiten theory we refer to [ASS] or [Ben1]. Let F be a field, B be an indecomposable Frobenius algebra over F with Nakayama automorphism ν and let P be a projective indecomposable B -module. We denote by τ the Auslander-Reiten translation of B . As B is a Frobenius algebra, we have $\tau \cong \Omega^2 \nu$ by [Ben1, 4.12.9]. Furthermore we denote by $B\text{-mod}$ the category of finite-dimensional left B -modules.

Let $\alpha \in \text{Aut}(B)$ and $M, N \in B\text{-mod}$. Then we denote by M_α the module which is isomorphic to M as an abelian group and the action of B on M_α is given by $b.m := \alpha(b)m$ for all $b \in B$ and $m \in M$. We denote by $l(M)$ the length of composition series of M , by $c(M)$ its complexity and by $\text{Irr}(M, N)$ the space of irreducible maps from M to N .

Let G be a subgroup of $\text{Aut}(B)$. We call B a G -transitive block, if for any two simple left B -modules V and W there is an element $g \in G$ such that $V_g \cong W$. We denote by $T_s(B)$ the stable Auslander-Reiten quiver of B .

From now on we assume that B is G -transitive.

We first prove a result on the length of the modules appearing at the end of an Auslander-Reiten sequence.

Lemma 2.1. *Let M be an indecomposable, non-projective B -module. Then $l(\tau(M)) = l(M) + nl(P)$ for some $n \in \mathbb{Z}$.*

Proof. As B is G -transitive all projective indecomposable modules of B have the same length. Therefore the length of projective covers of left B -modules are multiples of $l(P)$. It follows that for all $i \in \mathbb{N}$ there is a $n \in \mathbb{Z}$ such that $l(\Omega^i(M)) = l(M) + nl(P)$. Therefore $l(\tau(M)) = l(\Omega^2(M_{\nu^{-1}})) = l(M_{\nu^{-1}}) + nl(P) = l(M) + nl(P)$ for some $n \in \mathbb{Z}$. $\square$

The next lemma proves a condition which ensures that all indecomposable length 2 modules of a block have complexity one.

Lemma 2.2. *Suppose P has length 4. Then every indecomposable B -module of length 2 has complexity one.*

Proof. Let M be an indecomposable B -module of length 2. The module M has a simple top and therefore an indecomposable projective cover. Then $\Omega(M)$ is also an indecomposable length 2 module and has therefore an indecomposable projective cover. It follows that $l(\Omega^n(M)) = 2$ for all $n \in \mathbb{N}$. Therefore the complexity of M is one. $\square$

In general not every module of a Frobenius algebra with complexity 1 is periodic. In particular it does not need to hold for an algebra with Auslander-Reiten component of Euclidean tree class, as is shown in the next

Example 2.3. [SM] Let $A_q := F\langle x, y \rangle / (x^2, y^2, xy - qyx)$ where $q \neq 0$ and where q is not a root of unity. Let $M_\gamma = \text{Span}\{v, xv\}$ be the two-dimensional module with $yv = \gamma xv$ for $\gamma \in F^*$. The projective cover of M_γ is given by $\pi : A_q \rightarrow M_\gamma$ with $\pi(1) = v$. Then $\Omega(M_\gamma) = \text{Span}\{xy, y - \gamma x\} \cong M_{\gamma q}$. As q is not a root of unity we have $\Omega^k(M_\gamma) \not\cong M_\gamma$ for all $k \in \mathbb{N}$. Therefore M_γ is not periodic but has complexity one. Furthermore the Auslander-Reiten component containing the simple module is isomorphic to $\mathbb{Z}[\tilde{A}_{12}]$.

In the case of Auslander-Reiten components with Euclidean tree class, we know that the simple modules are non-periodic.

Lemma 2.4. *Suppose that $T_s(B)$ has a component θ of Euclidean tree class. Then all simple modules are non-periodic and lie in an Auslander-Reiten component isomorphic to θ .*

Proof. As θ has Euclidean tree class it is attached to a projective indecomposable module P by [W, 2.4] and [ASS, IV, 5.5]. If $l(P) < 4$, then $l(P)$ is uniserial which is a contradiction to the Euclidean tree class by [Ben1,

4.16.2]. As P is attached to θ the indecomposable module $P/\text{soc } P$ lies in θ . The map induced by Ω restricted to θ is an isomorphism. Therefore $\Omega(P/\text{soc } P) = \text{soc } P$ is contained in a component isomorphic to θ . So $\text{soc}(P)$ is not periodic. As B is G -transitive, all simple modules are non-periodic and lie in components isomorphic to θ . $\square$

We define certain stable graph automorphisms for G -transitive blocks with an Auslander-Reiten component θ of Euclidean tree class. Note that by [W, 2.4] and [ASS, IV, 5.5], there is at least one projective indecomposable module P attached to θ . Those maps have been defined in the proof of [F2, 4.6] similarly.

Definition 2.5. *Suppose that the stable Auslander-Reiten quiver $T_s(B)$ has a component θ of Euclidean tree class. Define $\phi_g : T_s(B) \rightarrow T_s(B)$ by $M \mapsto \Omega(M_g)$. Then ϕ_g is a stable graph isomorphism. Define for any $g \in G$ the map $A_g : T_s(B) \rightarrow T_s(B)$, by $M \mapsto M_g$. We denote by θ_g the component of $T_s(B)$ which is the image of $A_g(\theta)$.*

For the rest of the section we fix for any component θ with Euclidean tree class a projective indecomposable module P that is attached to θ and an element $g \in G$ such that $\phi := \phi_g|_\theta$ is an automorphism of θ . Furthermore let $S := \phi(P/\text{soc } P)$.

Also $\phi_g|_\theta$ is an automorphism of θ if and only if $P_{g^{-1}}$ is attached to $\Omega(\theta)$. We can see this as follows. If $P_{g^{-1}}$ is attached to $\Omega(\theta)$, then A_g induces an isomorphism from $\Omega(\theta)$ to θ and $\phi_g|_\theta$ is an automorphism.

Suppose $\phi_g|_\theta$ is an automorphism. Then $A_{g^{-1}}$ induces an isomorphism from θ to $\Omega(\theta)$. As P is attached to θ , $P_{g^{-1}}$ is attached to $\Omega(\theta)$.

Note that S is a simple module that belongs to θ .

First we need to show that the twisting action of ν commutes with the twisting action of any automorphism of B .

Lemma 2.6. *Let A be a Frobenius algebra. For all $g \in \text{Aut}(A)$ and A -modules M we have $M_{\nu g} \cong M_{g\nu}$.*

Proof. Let $(-, -)$ and $\{-, -\}$ be two associative non-degenerate bilinear forms, ν and ν_1 the corresponding Nakayama automorphisms and let $f := (-, 1)$ and $f_1 := \{-, 1\}$ be the corresponding linear forms. Then $\pi : A \rightarrow A^*, a \mapsto af$, and $\pi_1 : A \rightarrow A^*, a \mapsto af_1$, are B -module isomorphisms. Therefore there exist $x, y \in A$ such that $xf = f_1$ and $yf_1 = f$. It follows that $x = y^{-1}$. Set $u := \nu(x)$. Then the following equation holds: $\{a, b\} = (ab, x) = (a, bx) = (\nu(b)u, a) = \{\nu(b)u, ax^{-1}\} = \{u^{-1}\nu(b)u, a\}$ for all $a, b \in A$. Let $C_u : A \rightarrow A, a \mapsto u^{-1}au$ for all $a \in A$. Then $\nu_1 = C_u \circ \nu$.

Let $g \in \text{Aut}(A)$, then $\{-, -\} := (-, -) \circ (g \times g)$ is a associative non-degenerate bilinear form. It has Nakayama automorphism $g^{-1} \circ \nu \circ g$ as the following holds: $\{a, b\} = (g(a), g(b)) = (\nu(g(b)), g(a)) = \{g^{-1}(\nu(g(b))), a\}$

for all $a, b \in A$. By the first part there exists an invertible element $u \in A$ such that $g^{-1} \circ \nu \circ g = C_u \circ \nu$. Therefore $M_{g\nu g^{-1}} \cong M_{C_u \circ \nu}$ for all left B -modules M . But $M_\nu \cong M_{C_u \circ \nu}$ via the automorphism $\phi : M_\nu \rightarrow M_{C_u \circ \nu}, m \mapsto u^{-1}m$ and therefore we conclude $M_{g\nu} \cong M_{\nu g}$. $\square$

3. RESTRICTIONS ON TREE CLASSES

We say for the rest of the article that the algebra A satisfies (C) if

Every module with complexity 1 is Ω -periodic

We say an algebra satisfies (C') if

Every module with complexity 1 is Ω and τ -periodic

Note that if ν has finite order and (C) holds then (C') is also true. We introduce the following

Assumption 3.1. *For B , we assume that all elements in G have finite order. Furthermore we assume that B satisfies (C') and that $T_s(B)$ has a component θ of Euclidean tree class. Let P be a projective indecomposable module attached to θ .*

We have the following condition for the existence of a non-periodic indecomposable module of length 3.

Lemma 3.2. *Suppose B satisfies (C). If all indecomposable modules of length 2 have complexity one, then all indecomposable modules of length 3 are non-periodic. Also there is no uniserial module of length 3.*

Proof. Every indecomposable module of length 3 has a simple top or a simple socle. Let M be an indecomposable module of length 3 with simple top. Such an element exists, because a factor module of P of length 3 has simple top and is therefore indecomposable. Then there exists an exact sequence $0 \rightarrow S \rightarrow M \rightarrow L \rightarrow 0$ such that L is an indecomposable module of length 2 and S is a simple module. Then $c(S) \leq \max\{c(M), c(L)\}$. By 2.4 S is non-periodic and therefore $c(S) \geq 2$. As $c(L) = 1$, we have $c(M) \geq 2$ and therefore M has to be non-periodic. If M has a simple socle we can find an indecomposable module N of length 2 such that $0 \rightarrow N \rightarrow L \rightarrow S \rightarrow 0$ is an exact sequence. By the same argument as in the previous case, M has to be non-periodic.

We assume that there is a uniserial module M of length 3 with composition series S_1, S_2, S_3 . Let $L = \text{rad } M$ and $N = M/S_3$. Then an exact sequence is given by $0 \rightarrow L \rightarrow S_2 \oplus M \rightarrow N \rightarrow 0$. As L and N are indecomposable modules of length 2 they are periodic. Therefore $c(M) \leq 1$ and M is therefore periodic. This is a contradiction to the first part. $\square$

The proof of the next Theorem goes along the lines of the proof of [F2, 4.6]. As the setup here is more general and the author uses properties of the universal enveloping algebra of restricted p -Lie algebras, we give a proof for our setup.


Theorem 3.3. *Let B be as in 3.1. Then the following statements hold:*

- (1) θ is isomorphic to $\mathbb{Z}[\tilde{A}_{12}]$ or $\mathbb{Z}[\tilde{D}_n]$ with n odd.
- (2) The group G acts transitively on the non-periodic components.
- (3) If $\theta \cong \mathbb{Z}[\tilde{A}_{12}]$, then all projective indecomposable left B -modules have length 4. Furthermore all indecomposable modules of length $2 \bmod 4$ and all indecomposable non-projective modules of length $0 \bmod 4$ are periodic.

Proof. Let T be the tree class of θ . Then ϕ induces a graph automorphism $f : T \rightarrow T$. Suppose f has a fixed point. Then there is an indecomposable module M in θ such that $\Omega^{2m}(M_{\nu^{-m}}) \cong \tau^m(M) \cong \Omega(M_g)$. Therefore M has complexity 1 and is by assumption τ -periodic which is a contradiction. Therefore f does not have fixed points. The only Euclidean trees which admit an automorphism without fixed points are $\tilde{A}_{12}$ and $(\tilde{D}_n)_{n \geq 4}$ with n odd. We have $\theta \cong \mathbb{Z}[\tilde{A}_{1,2}]$ or $\theta \cong \mathbb{Z}[\tilde{D}_n]$ by [F2, 2.1]. Furthermore all indecomposable modules which do not lie in $\Psi := \bigcup_{w \in G} \theta_w$ are periodic. This can be seen as follows: let Y be an indecomposable module which is not in Ψ . We have $\Omega(\Psi) = \Psi$. Therefore the function $d_Y : \Psi \rightarrow \mathbb{N}, M \mapsto \dim_F \text{Ext}^1(Y, M)$ is additive by [ES, 3.2] and bounded by [W, 2.4]. Since all simple modules are contained in Ψ by 2.4 there exists an $m \in \mathbb{N}$ such that $\dim_F \text{Ext}^n(Y, W) \leq m$ for all $n \geq 1$ and all simple modules W , so that Y has complexity one. Therefore Y is periodic.

To prove (3) we suppose that $T = \tilde{A}_{12}$. Then the proof is the same as in [F2, 4.6]. For the convenience of the reader we include a different proof. As S and $P/\text{soc } P$ are in $\theta \cong \mathbb{Z}[\tilde{A}_{1,2}]$, all modules in θ have length $-1 \bmod l(P)$ or $1 \bmod l(P)$ and two modules connected by an arrow have a different length $\bmod l(P)$. The length function $l \bmod l(P)$ is additive on Auslander-Reiten sequences. Therefore $-2 \bmod l(P) = 2$ and it follows that $l(P) = 4$. As no modules of length $2 \bmod l(P)$ and $0 \bmod l(P)$ occur in θ , we have by (2) that all indecomposable modules of length $2 \bmod l(P)$ and all indecomposable non-projective modules of length $0 \bmod l(P)$ are periodic. $\square$

Let $\mathbb{Z}[\tilde{D}_n]$ be indexed as follows: $(k, 1), \dots, (k, n+1)$ denote the k -th copy of $\tilde{D}_n$ for any $k \in \mathbb{Z}$:



With this notation $\tau((k, i)) = (k-1, i)$.


For an indecomposable non-projective module we denote by $\bar{\alpha}(M)$ the number of predecessors of M in $T_s(B)$.

In order to prove our second Theorem, we need the following

Lemma 3.4. *Assume $T_s(B)$ has a component θ which is isomorphic to $\mathbb{Z}[\tilde{D}_n]$ with n odd. Then*

- (a) *$\text{rad}P/\text{soc}P$ is indecomposable.*
- (b) *$l(P)$ is even.*
- (c) *All M with $\bar{\alpha}(M) = 2$ or 3 have even length.*

Proof. (1) We assume that $l(P)$ is even and show that (c) holds in this case. Let M be a module with $\bar{\alpha}(M) = 3$. Then there exist a module N such that M is its only non-projective predecessor. Let $l(N) = a$, then $l(M) = 2a \bmod l(P)$ by Lemma 2.1. By assumption $l(P)$ is even and therefore all modules with 3 predecessors have even length. Consider the following extract from $\mathbb{Z}[\tilde{D}_n]$ where $M_3 := (k, 3)$ and $M_{n-1} := (k, n-1)$ denote the isomorphism type of modules with 3 predecessors and $M_t := (k, t)$ for $3 \leq t \leq n-1$:



We assume (c) is false and let t be minimal such that M_t has odd length. Suppose $t > 4$, then M_{t-2} is of even length and $\bar{\alpha}(M_{t-1}) = 2$. Then

$2l(M_{t-1}) = l(M_t) + l(M_{t-2}) \bmod l(P)$. This gives a contradiction as the right hand side is an even number and the left hand side is an odd number. Therefore $t = 4$. Then M_5 is of even length and M_6 of odd length. We can show that M_s with s odd has even length and M_s with s even has odd length. As n is odd, the module M_{n-1} has to be of odd length, which is a contradiction.

(2) The Auslander-Reiten sequence ending in $P/\text{soc } P$ is given by

$$0 \rightarrow \text{rad } P \rightarrow P \oplus \text{rad } P/\text{soc } P \rightarrow P/\text{soc } P \rightarrow 0.$$

We assume that $\text{rad } P/\text{soc } P$ is decomposable.

Then $\bar{\alpha}(P/\text{soc } P) > 1$. There exists no module N such that $0 \rightarrow \tau(N) \rightarrow S \rightarrow N \rightarrow 0$ is an Auslander-Reiten sequence. Any projective indecomposable module Q appears only as a summand of the middle term of an Auslander-Reiten sequence with middle term $Q \oplus \text{rad } Q/\text{soc } Q$ and $S \neq \text{rad } Q/\text{soc } Q$ because $l(Q) > 3$. So there exists no Auslander-Reiten sequence of the form $0 \rightarrow \tau(N) \rightarrow S \oplus Q \rightarrow N \rightarrow 0$ with Q non-zero. Therefore $\bar{\alpha}(S) \neq 3$. As ϕ maps $P/\text{soc } P$ to S we have $\bar{\alpha}(S) = \bar{\alpha}(P/\text{soc } P)$. Therefore $\bar{\alpha}(S) = \bar{\alpha}(P/\text{soc } P) = 2$. For $n = 5$ all modules have either one or three predecessors, so that $n > 5$. Suppose $l(P)$ is even. Then (1) shows that all modules with 2 or 3 predecessors in θ are of even length. As S is not of even length, this is a contradiction. Therefore $l(P)$ is odd. Let M_1 be a module of length a with only one predecessor M_3 , then $l(M_3) = 2a \bmod l(P)$. For the other module M_2 with only predecessor M_3 and length $\bar{a}$ we have $2\bar{a} = 2a \bmod l(P)$. As $l(P)$ is odd this gives us $\bar{a} = a \bmod l(P)$. We can deduce that $l(M_4) + 2a = 4a \bmod l(P)$ and therefore $l(M_4) = 2a \bmod l(P)$. It follows inductively that $l(M_i) = 2a \bmod l(P)$ for all modules with 2 predecessors. Therefore we have $-1 = 2a \bmod l(P)$ and $1 = 2a \bmod l(P)$ as $P/\text{soc } P$ and S are modules with 2 predecessors. This is a contradiction as $l(P)$ is odd. This proves (a).

(3) The only predecessor of $P/\text{soc } P$ is $\text{rad } P/\text{soc } P$. As $S = \phi(P/\text{soc } P)$, we have $\bar{\alpha}(S) = 1$. The predecessor of S has length $2 \bmod l(P)$ and the predecessor of $P/\text{soc } P$ has length $-2 \bmod l(P)$ by the argument of (2).

Suppose that $l(P)$ is odd. If $n = 5$ this is a contradiction. If $n > 5$, then by (2) all modules with 2 predecessors have length $2 \bmod l(P)$ and $-2 \bmod l(P)$ which is a contradiction as $l(P)$ is odd. Therefore (b) holds. Then (1) proves (c). $\square$

We also need the following

Lemma 3.5. *Suppose $T_s(B)$ has a component θ of tree class $\tilde{D}_n$. Then*

(1) $l(P) > 4$.

(2) $P/\text{soc } P$ and S have one predecessor and the τ -orbit of their predecessors are different.

Proof. We set $l := l(P)$. We know by Lemma 3.4 that $\text{rad } P/\text{soc } P$ is indecomposable.


It was shown in the proof of 2.4 that $l \geq 4$. Suppose now that $l = 4$, then by Lemma 2.2 all indecomposable modules of length 2 are periodic. By 3.4 $\text{rad } P/\text{soc } P$ is indecomposable and therefore periodic. As P is attached to θ , the module $\text{rad } P/\text{soc } P$ also belongs to θ which is a contradiction by [Ben1, 4.16.2]. Therefore $l > 4$.

From (3) of the proof of 3.4 we know that S and $P/\text{soc } P$ have only one predecessor of length $2 \bmod l$ and $-2 \bmod l$ respectively. As $l \neq 4$ and by 2.1 their predecessors do not lie in the same τ -orbit. □

We can now deduce the length of projective indecomposable modules if the Auslander-Reiten quiver has components $\mathbb{Z}[\tilde{D}_5]$.

Proposition 3.6. *Let B be as in 3.1. Suppose $T_s(B)$ has a component θ of tree class $\tilde{D}_5$. Then $l(P) = 8$ and all indecomposable modules of length 2 and of length $4 \bmod l(P)$ are periodic.*

Proof. We set $l := l(P)$. Let x be the length modulo l of the module which has as only predecessor the module of length $-2 \bmod l$ and let y the length modulo l of the module which has as only predecessor the module of length $2 \bmod l$. We visualize this in the following diagram:



Then by comparing lengths in Auslander-Reiten sequences we get the following equations:

- (1) $2x + 2 = 0 \bmod l$
- (2) $x + 5 = 0 \bmod l$
- (3) $5 - y = 0 \bmod l$
- (4) $2y - 2 = 0 \bmod l$

We can therefore deduce from (1) and (2) that l divides 8. Therefore $l = 8$ by 3.5 part (1).

Suppose now that there is an indecomposable module of length 2 which is not periodic. By transitivity there is an indecomposable non-periodic module M of length two in θ . Then by the equations (1)-(4), $\bar{\alpha}(M) = 3$ and there is an indecomposable module N which has only M as predecessor. Therefore M appears in the Auslander-Reiten sequence $0 \rightarrow \tau(N) \rightarrow M \rightarrow N \rightarrow 0$. This means that N and $\tau(N)$ have length one and are simple modules. By

transitivity $\tau(Q)$ is a simple B -module for any simple B -module Q . Then N has to be periodic which is a contradiction. By the equations (1)-(4), θ does not have an indecomposable module of length 4 mod l . Therefore those modules are periodic by 3.3. $\square$

We can now exclude tree class $\tilde{D}_5$ for certain algebras.

Theorem 3.7. *Let B be as in 3.1. Then $T_s(B)$ does not have a component with tree class $\tilde{D}_5$.*

Proof. We assume, for a contradiction, that $T_s(B)$ has a component θ of tree class $\tilde{D}_5$. Using 3.6 and 3.2, we know that B does not have a uniserial module of length 3, but has a non-periodic indecomposable module of length 3. Therefore x in the proof of 3.6 is 3. By the proof of 3.6 there is an almost split sequence $0 \rightarrow \tau(X) \xrightarrow{f} H \xrightarrow{g} X \rightarrow 0$ with $H := \text{rad } P / \text{soc } P$. Therefore $l(\tau(X)) = l(X) = 3$. Suppose $\tau(X)$ has an indecomposable submodule U of length 2. Then H has also an indecomposable submodule $V := f(U)$ of length 2. But then the preimage of V of the canonical surjection $\text{rad}(P) \rightarrow H$ is a submodule of length 3 and is uniserial which is a contradiction. Therefore $\tau(X)$ has a quotient W that is indecomposable of length 2. Let $h : \tau(X) \rightarrow W$ be the canonical surjection. Then by Auslander-Reiten theory h factors through f . Therefore there exists a surjective map $s : H \rightarrow W$. But then $P / \ker s$ is a uniserial module of length 3 which is a contradiction. Therefore $T_s(B)$ does not have a component of tree class $\tilde{D}_5$. $\square$

We define the following automorphisms of $\mathbb{Z}[\tilde{D}_n]$ as in [F2].

$$\begin{aligned} \alpha(k, i) &= \begin{cases} (k, 1), & i = 2 \\ (k, 2), & i = 1 \\ (k, i), & 3 \leq i \end{cases} \\ \beta(k, i) &= \begin{cases} (k, n+1), & i = n \\ (k, n), & i = n+1 \\ (k, i), & i \leq n-1 \end{cases} \\ \gamma(k, i) &= \begin{cases} (k, n), & i = 1 \\ (k, n+1), & i = 2 \\ (k+i-3, n+2-i), & 3 \leq i \leq n-1 \\ (k+n-4, 1), & i = n \\ (k+n-4, 2), & i = n+1 \end{cases} \end{aligned}$$

Lemma 3.8. [F2, 2.1] *The automorphism group of $\mathbb{Z}[\tilde{D}_n]$ is given by*

$$\{\tau^k \circ \alpha^i \circ \beta^j \circ \gamma^l \mid k \in \mathbb{Z}, i, j, l \in \{0, 1\}\}.$$

We describe the action of G on Euclidean components.

Lemma 3.9. *Let B be as in 3.1. Let $h \in G$ and suppose h induces an automorphism $A_h : \theta \rightarrow \theta, M \mapsto M_h$. Suppose that B has an indecomposable*

non-periodic module of length 3, if θ has tree class $\tilde{D}_n$ for $n > 5$. Then A_h is the identity.

Proof. By 3.3 we have that $\theta \cong \mathbb{Z}[\tilde{D}_n]$ or $\theta \cong \mathbb{Z}[\tilde{A}_{1,2}]$. We assume first that $\theta \cong \mathbb{Z}[\tilde{D}_n]$ with $n > 5$. Suppose A_h is not the identity. By 3.8 the automorphisms of finite order have the form $\tau^k \circ \alpha^i \circ \beta^j \circ \gamma$ for $k = n/2 - 2$ or $\alpha^i \circ \beta^j$ with $i, j \in \{0, 1\}$. As n is odd the first possibility cannot occur. Therefore A_h is equal to either α , β or $\alpha \circ \beta$.

Suppose the map A_h is equal to $\alpha \circ \beta$. Then all modules with only one predecessors have length $\pm 1 \pmod{l(P)}$. There exists a non-periodic indecomposable module of length 3 and by transitivity there is an indecomposable length 3 module M in θ . As $l \neq 4$ by 3.5 part (1) we have $\bar{\alpha}(M) = 3$ or $\bar{\alpha}(M) = 2$. Therefore $M_h \cong M$. This is a contradiction because M has either a simple top or a simple radical and the map A_h does not stabilize simple modules.

Assume that $A_h = \beta$. Then $A_h(P/\text{soc } P) = P_h/\text{soc } P_h \not\cong P/\text{soc } P$. By definition of S and ϕ , we have $S = \text{soc } P_g$. Then $S = \text{soc } P_g = \phi(P/\text{soc } P) \not\cong \phi(P_h/\text{soc } P_h) = \text{soc } P_{hg} = S_{g^{-1}hg}$. Therefore $A_{g^{-1}hg} = \alpha$ as by the first case no automorphism induced by an element of G is equal to $\alpha \circ \beta$. But then $A_{hg^{-1}hg} = \alpha \circ \beta$, which is a contradiction.

Assume now that $A_h = \alpha$. Then $A_h(S) = S_h \not\cong S$. We have therefore $P/\text{soc } P = \phi^{-1}(S) \not\cong \phi^{-1}(S_h) = P_{ghg^{-1}}/\text{soc } P_{ghg^{-1}}$. Therefore $A_{ghg^{-1}} = \beta$ and $A_{hghg^{-1}} = \alpha \circ \beta$, which is a contradiction.

In the case of $\mathbb{Z}[\tilde{A}_{1,2}]$ there are no finite order automorphisms unequal to the identity, so this gives a contradiction as well. $\square$

We describe the non-periodic components more precisely in the following

Corollary 3.10. *Let B be as in 3.9. Then B has exactly as many non-periodic Auslander-Reiten components as there are isomorphism classes of simple left B -modules.*

Proof. By Theorem 3.3 all non-periodic components are isomorphic and for every non-periodic Auslander-Reiten component Δ there exists a $g \in G$ such that $\theta_g = \Delta$. The component θ contains a simple module by 2.4 and therefore every non-periodic Auslander-Reiten component contains a simple module. By transitivity there exists for any simple module V a non-periodic Auslander-Reiten component $\mathcal{W}$ such that V belongs to $\mathcal{W}$. Suppose there is a non-periodic component which contains two simple modules V and V_r for some $r \in G$. Then r induces a non-identity automorphism of finite order on the component. This is a contradiction to 3.9. $\square$

We can now prove some necessary conditions for a component of tree class $\tilde{D}_n$ for $n > 5$. Compare the following Theorem to [F2, 4.6]. We have proved

that $n \neq 5$. Farnsteiner first shows that $n = 5$ and then deduces the other statements from this fact. As this step is wrong, we require a different proof.

Theorem 3.11. *Let B be as in 3.1. Suppose $T_s(B)$ has a component θ of tree class $\tilde{D}_n$, $n > 5$. Suppose B contains an indecomposable non-periodic module of length 3. Then $l(P) = 8$ and all modules of length $4 \bmod l(P)$ are periodic.*

Proof. The proof follows in two steps.

Let $l := l(P)$.

Step 1: $l = 8$.

Let M be an indecomposable length 3 module in θ . By (c) of 3.4, $\bar{\alpha}(M) = 1$. Suppose M shares a predecessor with the module of length $1 \bmod l$. Then the predecessor has length $2 \bmod l$ and $6 \bmod l = 2 \bmod l$ which is a contradiction as $l \neq 4$. It must therefore share a predecessor with the module of length $-1 \bmod l$. This gives us $6 = -2 \bmod l$ and therefore $l = 8$.

We know from 3.5 (2) that the modules with 3 predecessors have length $2 \bmod 8$ and $-2 \bmod 8$. The modules with one predecessor have therefore length $\pm 1 \bmod 8$ or $\pm 3 \bmod 8$.

Step 2: The indecomposable modules of length $4 \bmod 8$ are periodic.

Let W be a module with one predecessor and length $1 \bmod 8$. We take W corresponding to $(k, 1)$ and use the notation of the proof 3.4. Then $l(M_3) = 2 \bmod 8$. Let $\bar{W}$ be the other module with only predecessor M_3 . Then $l(\bar{W}) = 4x + 1$. The module $l(M_4)$ satisfies $1 + l(\bar{W}) + l(M_4) = 4 \bmod 8$. Therefore $l(M_4) = 2(1 - 2x) \bmod 8$. In the same way we follow $l(M_5) = 2 \bmod 8$, $l(M_6) = 2(1 + 2x) \bmod 8$, $l(M_7) = 2 \bmod l$, $l(M_8) = 2(1 - 2x) \bmod 8$. The calculation shows that $l(M_t) = 2 \bmod 8$ if t is odd, $l(M_t) = 2(1 + 2x) \bmod 8$ if $t = 4m + 2$ and $l(M_t) = 2(1 - 2x) \bmod 8$ if $t = 4m$ for any $m \in \mathbb{N}$.

Thus modules of length $4 \bmod 8$ in θ do not have two predecessors. By the remark before Step 2 they do not have one or three predecessors. Therefore no module of length $4 \bmod 8$ belongs to θ . As no module of length $4 \bmod 8$ appears in θ , they have to be periodic by (2) of 3.3. $\square$

Note also that by the proof of 3.7 B has a uniserial module of length 3.

4. APPLICATION TO AUSLANDER-REITEN COMPONENTS OF ENVELOPING ALGEBRAS OF RESTRICTED p LIE ALGEBRAS

Let L be a finite-dimensional restricted p -Lie algebra and χ a linear form on L . We denote by $u(L, \chi)$ the universal enveloping of (L, χ) . If $\chi = 0$ we set $u(L, \chi) = u(L)$.

We denote by $G(L)$ the set of group-like elements of the dual Hopf algebra $u(L)^*$. The set of group-like elements are the homomorphisms of $u(L)$. The comultiplication on $u(L)$ induces an algebra homomorphism $\Delta : u(L, \chi) \rightarrow u(L) \otimes u(L, \chi), x \mapsto x \otimes 1 + 1 \otimes x$ for all $x \in L$. We denote $\Delta(u) = u_1 \otimes u_2$ for $u \in u(L, \chi)$. This defines a left $u(L)$ -comodule algebra structure and right $u(L)^*$ -module algebra structure on $u(L, \chi)$. Therefore $G(L)$ acts on the automorphism group of $u(L, \chi)$ via $(g \cdot \psi)(u) = \psi(u \cdot g) = g(u_1)\psi(u_2)$ for all $\psi \in \text{Aut}(u(L, \chi))$, $g \in G(L)$ and $u \in u(L, \chi)$. We embed $G(L)$ into $\text{Aut}(u(L, \chi))$ via the injective group homomorphism $f : G(L) \rightarrow \text{Aut}(u(L, \chi)), w \mapsto w \cdot \text{id}_{u(L, \chi)}$. For an $u(L, \chi)$ -module M and $w \in G(L)$ we denote by M_w the twisted module $M_{f(w)}$. Note that every element of $G(L) \setminus \{1\}$ has order p .

By [FS1, 1.2] the Nakayama automorphism of $u(L, \chi)$ has order 1 or p and all modules of complexity one are 2-periodic by [F1, 2.5]. Furthermore $u(L, \chi)$ has a non-periodic indecomposable module of order 3 by [F2, 4.5]. Therefore the assumptions of 3.1 are satisfied for $G(L)$ -transitive blocks or G -transitive blocks, where G is a finite subgroup of $\text{Aut}(u(L, \chi))$. The next corollary follows directly from 3.7.

Corollary 4.1. *Let $B \subset u(L, \chi)$ be a G -transitive block, then $T_s(B)$ does not have a component of tree class $\tilde{D}_5$.*

More generally we have

Lemma 4.2. *Let $B \subset u(L, \chi)$ be a G -transitive block and let S be a simple module in B , then $T_s(B)$ admits an Euclidean component only if $p = 2$ or $\dim S = 0 \bmod p$.*

Proof. By 3.3 and 3.11 all indecomposable modules of length two or of length four are periodic. By [F1, 2.5] all periodic indecomposable modules have dimension $0 \bmod p$. As B is G -transitive all simple modules in B have the same dimension. Therefore $2 \dim S = 0 \bmod p$. □

As a G -transitive principal block of $u(L, \chi)$ has only one dimensional simples, we get the following corollary immediately from the preceding Lemma.

Corollary 4.3. *Let $B \subset u(L, \chi)$ be the principal block, and assume B is G -transitive, then $T_s(B)$ admits an Euclidean component only if $p = 2$.*

We call a block B primary if it only contains one isomorphism type of simple modules. Note that lemma [F2, 4.7] remains true for primary blocks of $u(L, \chi)$ with the additional assumption that all indecomposable modules of length 2 are periodic.

We remind of the definition of supersolvable Lie algebras

Definition 4.4. [FS, I] Let $(L^i)_{i \in \mathbb{N}}$ with $L^i = [L^{i-1}, L]$ and $L^0 = L$ be a sequence of ideals in L . Then L is nilpotent if there is an $n \in \mathbb{N}$ such that $L^n = 0$. The sequence $(L^{(i)})_{i \in \mathbb{N}}$ with $L^{(i)} = [L^{(i-1)}, L^{(i-1)}]$ and $L^{(0)} = L$ is the derived series. We call L solvable if there is an $n \in \mathbb{N}$ such that $L^{(n)} = 0$ and L supersolvable if L^1 is nilpotent.

Using the fact that projective modules of restricted universal enveloping algebras of supersolvable Lie algebras have p -power length by [F3, 2.10], the result of [F2, 4.1] remains true by applying 3.11.

Lemma 4.5. Let L be a supersolvable finite-dimensional restricted p -Lie algebra and $p > 2$. Then $T_s(u(L, \chi))$ does not have a component of Euclidean tree class.

This result cannot be extended to $p = 2$ as the following example shows.

Example 4.6. Let $A = k[x, y]/(x^2, y^2)$ be the Kronecker algebra. Then $A \cong u(L)$ where $L = \text{Span}\{x, y\}$ is the restricted 2-Lie algebra given by $[x, y] = 0$ and $x^{[2]} = y^{[2]} = 0$. Then L is supersolvable and the component containing the trivial module k is isomorphic to $\mathbb{Z}[\tilde{A}_{1,2}]$. This is well known, see for example [E, 2.3].

5. EUCLIDEAN COMPONENTS OF SMASH PRODUCTS

The goal of this section is to determine conditions so that the smash product of a basic simple algebra and a semi-simple commutative group algebra have an Auslander-Reiten component of Euclidean tree class. We assume that k is algebraically closed.

We start by describing the simple and indecomposable projective modules of certain smash products.

Lemma 5.1. Let Γ be a local and basic algebra with simple module S and let G be a finite group such that $G < \text{Aut}(\Gamma)$. Let $\{e_1, \dots, e_m\}$ be a full set of primitive orthogonal idempotents in kG , let $\bigoplus_{i=1}^m P_i$ be a decomposition of kG into projective indecomposable kG -modules $P_i := kGe_i$ and let $S_i := \text{soc } P_i$ for $1 \leq i \leq m$.

Then for every simple $\Gamma \rtimes kG$ -module V there exists an $1 \leq i \leq m$ such that $V \cong S \otimes S_i$. A complete set of primitive orthogonal idempotents of $\Gamma \rtimes kG$ is given by $\{1 \rtimes e_i | 1 \leq i \leq m\}$ and $\Gamma \rtimes kG$ has a decomposition $\bigoplus_{i=1}^m \Gamma \rtimes P_i$ into projective indecomposable modules $\Gamma \rtimes P_i$.

Proof. As g induces an automorphism on Γ for all $g \in G$, we have $G(J(\Gamma)) = J(\Gamma)$ and $J(kG)\Gamma \subset J(\Gamma)$. Therefore $J(\Gamma) \rtimes kG + \Gamma \rtimes J(kG) \subset J(\Gamma \rtimes kG)$. Furthermore $\Gamma \rtimes kG / (J(\Gamma) \rtimes kG + \Gamma \rtimes J(kG)) \cong \Gamma / J(\Gamma) \otimes kG / J(kG) \cong \bigoplus_{i=1}^m S \otimes S_i$ which is semi-simple. This proves $J(\Gamma) \rtimes kG + \Gamma \rtimes J(kG) = J(\Gamma \rtimes kG)$ and all simples are given by $S \otimes S_i$. Clearly $\{1 \rtimes e_i | 1 \leq i \leq m\}$

is a set of orthogonal idempotents and $\bigoplus_{i=1}^m \Gamma \rtimes P_i$ is a decomposition of $\Gamma \rtimes kG$ into projective modules $\Gamma \rtimes P_i = (\Gamma \rtimes kG)(1 \rtimes e_i)$. The projective modules are indecomposable as $\text{soc}(\Gamma \rtimes P_i) = S \otimes S_i$ is simple and therefore $\{1 \rtimes e_i | 1 \leq i \leq m\}$ is a complete set of primitive idempotents. $\square$

From now on let Γ be a basic local algebra with simple module S . Let G be an abelian group such kG is semi-simple, and G is a subgroup of $\text{Aut}(\Gamma)$. Then the smash product $R := \Gamma \rtimes kG$ is well defined.

By Gabriel's lemma [Ben1, 4.1.7], there exists a quiver Q such that $\Gamma \cong kQ/I$ for an admissible ideal $I \subset kQ$. As G is abelian and kG semi-simple, the set of irreducible characters of kG forms a multiplicative group isomorphic to G . We index the characters by elements of G via a fixed isomorphism and index the primitive orthogonal idempotents by the same group element as its corresponding character. So let $\{\chi_g | g \in G\}$ be the set of irreducible characters and $\{e_g | g \in G\}$ the set of primitive orthogonal idempotents, such that $he_g = \chi_g(h)e_g$ for all $g, h \in G$. Suppose $G \leq \text{Aut}(\Gamma)$. Then kG acts on $J(\Gamma)$ and $J^2(\Gamma)$. As kG is semi-simple, $J(\Gamma)/J^2(\Gamma)$ split as a direct sum of one-dimensional kG -modules. Let $\alpha_1, \dots, \alpha_m$ be the simultaneous eigenvectors of the action of G on $J(\Gamma)/J^2(\Gamma)$. Let $\chi_{n_i}, n_i \in G$, $i = 1, \dots, m$ be the corresponding irreducible characters. By 5.1 we know that $\Gamma \rtimes kG$ is a basic algebra with projective indecomposable modules $\Gamma \rtimes ke_g$ for $g \in G$ which have simple quotients $S \rtimes ke_g$. We have the following presentation of $\Gamma \rtimes kG$. Take the quiver where vertices are labelled by $1 \rtimes e_g$ and where arrows are $\alpha_i \rtimes e_g$. Note that

$$(\alpha_i \rtimes e_h)(\alpha_j \rtimes e_g) = (\alpha_i \alpha_j) \rtimes \chi_{n_j g}(e_h)e_g = (\chi_{n_j g}, \chi_h)(\alpha_i \alpha_j \rtimes e_g)$$

where $(-, -)$ is the usual inner product of characters. Therefore the arrow $\alpha_i \rtimes e_g$ ends in $1 \rtimes e_g$ and starts in $1 \rtimes e_q$ with $q = gn_i$. We can obtain the relations that generate T via the relations that generate I in Γ .

Note that the construction of W coincides with the Mc Kay quiver (see [SSS, 2] for the definition) where $V := J(\Gamma)/J^2(\Gamma)$.

We will illustrate this construction on a small example.


Example 5.2. Let $\Gamma = k[x, y]/\langle x^2, y^2 \rangle$ the Kronecker algebra and let $G = \langle g \rangle$ be a cyclic group of order 3.

Then $\Gamma \cong kQ/I$ with

$$Q = x \begin{array}{c} \circ \\ \curvearrowright \end{array} \bullet \begin{array}{c} \circ \\ \curvearrowleft \end{array} y$$

and $I = \langle x^2, y^2, xy - yx \rangle$. The algebra Γ is a kG -module algebra via the action $gx = q^{-1}x$, $gy = qy$ and $gxy = xy$ for a primitive third root of unity q . We label the character χ with $\chi(g) = q$ as $\chi = \chi_g$. Then $n_x = g^2$ and $n_y = g$. Let e_1, e_g and e_{g^2} denote the primitive idempotents in G such that $ge_1 = e_1$, $ge_g = qe_g$ and $ge_{g^2} = q^{-1}e_{g^2}$. Then the primitive idempotents

are given by $1 \rtimes e_i$. We construct the quiver W as described in the previous example.



The relations are given by

$$T := \langle (y \rtimes e_i)(y \rtimes e_j), (x \rtimes e_i)(x \rtimes e_j), (x \rtimes e_{gj})(y \rtimes e_j) - (y \rtimes e_{g^2j})(x \rtimes e_j) \mid i, j \in \{1, g, g^2\} \rangle.$$

By the previous example $\Gamma \rtimes kG \cong kW/T$.

We have that R is G -transitive via $g(a \rtimes h) = a \rtimes \chi_{g^{-1}}(h)h$ or $g(a \rtimes e_h) = a \rtimes e_{gh}$ for all $a \in \Gamma$ and $g, h \in G$. With this action G is a subgroup of $\text{Aut}(R)$. Note that $1 \rtimes G \cong G$ is a subgroup of R and $k \rtimes G \cong kG$ is a subalgebra of R . We first define the following notation. Let C be an R -module, then C is a kG -module via $g \cdot c := (1 \rtimes g)c$ for all $c \in C$ and $g \in G$. We denote by C_g the R -module with $(a \rtimes h) * c := \chi_{g^{-1}}(h)(a \rtimes h)c$ for all $c \in C$, $g, h \in G$ and $a \in \Gamma$. If C is a Γ -module, we denote by C_g the module with $t * c := g(t)c$ for all $c \in C$ and $t \in \Gamma$.

If C is a Γ or an R -module, then we set $S(C) := \{g \in G \mid C_g \cong C\}$ with the respective action of G on R and on Γ -modules. Let $T(C)$ be a transversal of $G/S(C)$.

We determine how R -modules or Γ -modules decompose if restricted to R or respectively lifted to R .

Lemma 5.3. (1) Let M be an R -module. Then $(M_\Gamma)^R \cong \bigoplus_{g \in G} M_g$.

(2) Let N be a Γ -module, then $N_\Gamma^R \cong \bigoplus_{g \in G} N_g$ where N_g denotes the twist of N by the element $g \in \text{Aut}(\Gamma)$.

(3) Let M be an indecomposable R -module and N an indecomposable Γ -module such that $N \mid (M_\Gamma)$. Then $M_\Gamma = q \bigoplus_{g \in T(N)} N_g$, $N^R = n \bigoplus_{g \in T(M)} M_g$ and $qn|T(N)||T(M)| = |G|$ for some $n, m \in \mathbb{N}$.

Proof. (1) Let $\psi_g : M_g \rightarrow (M_\Gamma)^R, m \mapsto |G|^{-1} \sum_{l \in G} \chi_g(l)(1 \rtimes l \otimes l^{-1}m)$ and let $\phi_g : (M_\Gamma)^R \rightarrow M_g, r \otimes m \mapsto g(r)m$ for all $r \in R, m \in M$ and $g \in G$. Let

$a \in \Gamma$ and $h \in G$. Then

$$\begin{aligned}
\psi_g((a \rtimes h) * m) &= \psi_g(\chi_{g^{-1}}(h)(a \rtimes h)m) \\
&= |G|^{-1} \sum_{l \in G} \chi_{g^{-1}}(h) \chi_g(l) (1 \rtimes l \otimes l^{-1}(a \rtimes h)m) \\
&= |G|^{-1} \sum_{l \in G} \chi_{g^{-1}}(h) \chi_g(l) (1 \rtimes l) (l^{-1}(a) \rtimes 1 \otimes l^{-1}hm) \\
&= |G|^{-1} (a \rtimes 1) \sum_{l \in G} \chi_{g^{-1}}(h) \chi_g(l) (1 \rtimes l \otimes l^{-1}hm) \\
&= (a \rtimes h) |G|^{-1} \sum_{s \in G} \chi_g(s) (1 \rtimes s \otimes s^{-1}m) \\
&= (a \rtimes h) \psi_g(m),
\end{aligned}$$

by substituting $l^{-1}h = s^{-1}$, and therefore ψ_g is an R -module homomorphism.

It is clear that ϕ_g is an R -module homomorphisms and $\phi_g \circ \psi_g = \text{id}_{M_g}$. Let $\{m_1, \dots, m_n\}$ be a k -basis of M . A basis of $(M_\Gamma)^R$ is given by

$$\{(1 \rtimes l) \otimes m_i \mid \text{for } 1 \leq i \leq n \text{ and } l \in G\}.$$

Using this basis we have $\psi_g(m) = \psi_h(\bar{m})$ for some $m, \bar{m} \in M$ if and only if $\chi_g(l)m = \chi_h(l)\bar{m}$ for all $l \in G$. Therefore $\psi_g(M_g) \cap \psi_h(M_h) = 0$ for $g \neq h$. Finally by comparing dimensions we have $(M_\Gamma)^R \cong \bigoplus_{g \in G} M_g$.

(2) We have $N_\Gamma^R = \bigoplus_{g \in G} 1 \rtimes g \otimes N$. Furthermore $1 \rtimes g \otimes N \cong N_{g^{-1}}$ as Γ -module, which proves the statement.

(3) Suppose Q is an indecomposable module with $Q|(M_\Gamma)$, then Q^R and N^R are direct summands of $(M_\Gamma)^R = \bigoplus_{g \in G} M_g$. As $(M_g)_\Gamma \cong M_\Gamma$ for all $g \in G$, we have that $N|(Q_\Gamma^R) = \bigoplus_{g \in G} Q_g$. Therefore $Q \cong N_g$ for some $g \in G$. Furthermore $(M_\Gamma)_g \cong M_\Gamma$ via the Γ -module isomorphism $\psi : M_\Gamma \rightarrow (M_\Gamma)_g, m \mapsto gm$ for all $m \in M$ and $g \in G$. Therefore M_Γ is G -invariant. This proves the first identity.

By the first identity, we know that all indecomposable direct summands of N^R are isomorphic to M_g for some $g \in G$. Note that N^R is G -invariant via the R -module isomorphism $\phi : N^R \rightarrow (N^R)_g, r \otimes n \mapsto g(r) \otimes n$ for all $g \in G, r \in R$ and $n \in N$. This map is well defined as G acts on $\Gamma \rtimes 1 \subset R$ as the identity. Therefore the second identity holds.

Finally we compare the multiplicity of N as a direct summand of N_Γ^R . The first and second identity of (3) give a multiplicity of $n|T(M)|q$ and (2) gives multiplicity $|S(N)|$. $\square$

By standard arguments we deduce the next two lemmas.

Lemma 5.4. *Every R -module M is relatively Γ -projective.*

Proof. Suppose A, B are R -modules and $h : A \rightarrow B, f : M \rightarrow B$ are R -module homomorphisms. Suppose there is a Γ -module homomorphism $v :$

$M_\Gamma \rightarrow A$ such that $h \circ v = f$. Then $\bar{v} : M \rightarrow A, m \mapsto |G|^{-1} \sum_{g \in G} gv(g^{-1}m)$ is an R -module homomorphism. This can be seen as follows: let $t \in \Gamma$, $h \in G$, then

$$\begin{aligned}
\bar{v}((t \rtimes h)m) &= |G|^{-1} \sum_{g \in G} gv(g^{-1}(t \rtimes h)m) \\
&= |G|^{-1} \sum_{g \in G} gv(g^{-1}(t) \rtimes g^{-1}h)m) \\
&= |G|^{-1}(t \rtimes 1) \sum_{g \in G} gv(g^{-1}hm) \\
&= (t \rtimes h)|G|^{-1} \sum_{s \in G} sv(s^{-1}m) \\
&= (t \rtimes h)\bar{v}(m),
\end{aligned}$$

if we substitute $s^{-1} = g^{-1}h$. Furthermore $\bar{v}$ satisfies $h \circ \bar{v} = f$. $\square$

Lemma 5.5 (Frobenius reciprocity). *Let V be a Γ -module and M an R -module. Then there is a bijection of vector spaces between $\text{Hom}_\Gamma(V, M_\Gamma)$ and $\text{Hom}_R(V^R, M)$.*

Proof. The bijection is given by $\psi : \text{Hom}_\Gamma(V, M_\Gamma) \rightarrow \text{Hom}_R(V^R, M)$ where $\psi(f)(r \otimes v) = rf(v)$ and $\phi : \text{Hom}_R(V^R, M) \rightarrow \text{Hom}_\Gamma(V, M_\Gamma)$ with $\phi(g)(v) = g(1 \otimes v)$ for all $r \in R, v \in V, f \in \text{Hom}_\Gamma(V, M_\Gamma)$ and $g \in \text{Hom}_R(V^R, M)$. $\square$

Let $G(A)$ denote the free abelian group of an algebra A with free generators $[V_i]$, the representatives of the isomorphism classes of all indecomposable A -modules V_i . If $M = \oplus a_i V_i$ where the $a_i \geq 0$ then we write $[M] := \sum a_i [V_i]$. We denote by $(-, -)_A : G(A) \times G(A) \rightarrow k$ the bilinear form $\dim_k \text{Hom}_A(-, -)$.

Let $Q : 0 \rightarrow B \rightarrow C \rightarrow D \rightarrow 0$ be an exact sequence. Then we set $[[Q]] := [B] + [D] - [C] \in G(A)$. Let $A(V_i)$ denote the Auslander-Reiten sequence starting in V_i for V_i non-projective. Furthermore we set $X_i := [[A(V_i)]]$ for V_i non-projective and $X_i := [V_i] - [\text{rad}(V_i)]$ if V_i is projective.

The first part of the next Theorem is the general version of [BP, 3.4], that was only proven for group algebras.

Theorem 5.6. *Let A be a finite-dimensional algebra over an algebraically closed field k .*

(1) *We have that $([V_i], X_j) = \delta_{i,j}$. Therefore $(-, -)_A$ is non-degenerate. Furthermore $([V_i], [[E]]) \geq 0$ for any exact sequence E .*

(2) *Suppose $Q := 0 \rightarrow C \rightarrow B \rightarrow V_j \rightarrow 0$ is an exact non-split sequence with $[[Q]] \neq X_j$, then there is a V_i with $j \neq i$ such that $([V_i], [[Q]]) \geq 1$.*

Proof. (1) Take the almost split sequence

$$0 \rightarrow \tau(V_j) \xrightarrow{l} M_j \xrightarrow{s} V_j \rightarrow 0$$

this gives an exact sequence

$$0 \rightarrow \operatorname{Hom}_A(V_i, \tau V_j) \rightarrow \operatorname{Hom}_A(V_i, M_j) \xrightarrow{\psi} \operatorname{Hom}_A(V_i, V_j).$$

If $i \neq j$ then by the Auslander-Reiten property, the map ψ is onto and it follows that $([V_i], X_j) = 0$. If $i = j$ then by the Auslander-Reiten property, $\operatorname{Im}(\psi)$ is the radical of $\operatorname{End}(V_i)$. Therefore

$$([V_i], X_i) = (V_i, \tau(V_i)) + (V_i, V_i) - (V_i, M_i) = (V_i, V_i) - \dim \operatorname{Im}(\psi)$$

and this is equal to

$$\dim(\operatorname{End}(V_i)/\operatorname{rad}(\operatorname{End}(V_i))).$$

Since k is algebraically closed and V_i is indecomposable, this number is equal to 1.

Let $E := 0 \rightarrow S \rightarrow T \rightarrow U \rightarrow 0$ be an exact sequence. Then $0 \rightarrow \operatorname{Hom}_A(V_i, S) \rightarrow \operatorname{Hom}_A(V_i, T) \rightarrow \operatorname{Hom}_A(V_i, U)$ is exact. Therefore $([V_i], [[E]]) \geq 0$.

(2) Let $Q := 0 \rightarrow C \xrightarrow{\delta} B \xrightarrow{\sigma} V_j \rightarrow 0$ be an exact sequence. Suppose that $[[Q]] \neq [[A(V_j)]]$. Then we get the following commutative diagram

$$\begin{array}{ccccccccc} 0 & \longrightarrow & C & \xrightarrow{\delta} & B & \xrightarrow{\sigma} & V_j & \longrightarrow & 0 \\ & & \downarrow g & & \downarrow h & & \downarrow \operatorname{id} & & \\ 0 & \longrightarrow & \tau(V_j) & \xrightarrow{l} & M_j & \xrightarrow{s} & V_j & \longrightarrow & 0 \end{array}$$

where the existence of h follows from the Auslander-Reiten property since the map $\sigma : B \rightarrow V_j$ is non-split; and g is the restriction of h to C .

This diagram induces a short exact sequence

$$Z := 0 \rightarrow C \xrightarrow{\left(\begin{smallmatrix} \delta \\ g \end{smallmatrix}\right)} B \oplus \tau(V_j) \xrightarrow{(h, l)} M_j \rightarrow 0.$$

Suppose that this sequence is split. Then there is a map $\begin{pmatrix} f_1 \\ f_2 \end{pmatrix} : M_j \rightarrow B \oplus \tau(V_j)$ such that $h \circ f_1 + l \circ f_2 = \operatorname{id}_{M_j}$. Then $s \circ h f_1 = s$. As s is minimal right almost split, we have that $h f_1$ is an automorphism. We also have $\sigma f_1 = s$, so let $g_1 : \tau(V_j) \rightarrow C$ such that $\delta g_1 = f_1 l$. Then also $g g_1$ is an automorphism. So we have $B = M_j \oplus \ker(h)$ and $C = \tau(V_j) \oplus \ker(g)$. By the Snake Lemma, $\operatorname{Ker}(g) \cong \operatorname{Ker}(h)$. But then $[[Q]] = [[A(V_j)]]$. So we have a contradiction, therefore Z is non-split. Then we have, working in $G(A)$, that $[[Q]] = [[Z]] + [[A(V_j)]]$ and hence we have $([M_j], [[Q]]) = ([M_j], [[Z]] + [[A(V_j)]]) = ([M_j], [[Z]]) \geq 1$ as the image of the map $\operatorname{Hom}(M_j, M_j) \rightarrow \operatorname{Ext}^1(M_j, C)$ induced by Z contains the non-split sequence Z . As $(-, -)_A$ is biadditive and V_j is not a summand of M_j , the second statement is proven. $\square$

We can write the element $[[Q]] \in G(A)$ for any exact sequence Q ending in W as a sum in $G(A)$ of $[[Q_1]]$ and $[[Q_2]]$ for two short exact sequences Q_1 and Q_2 ending in direct summands of W .

Lemma 5.7. *Suppose that $Q := 0 \rightarrow U \rightarrow V \xrightarrow{\pi} W \rightarrow 0$ is an exact sequence and $W = W_1 \oplus W_2$ for two non-trivial A -modules W_1 and W_2 . Then there is an exact sequence Q_1 ending in W_1 and an exact sequence Q_2 ending in W_2 such that $[[Q]] = [[Q_1]] + [[Q_2]]$.*

Proof. Let $p_i : W \rightarrow W_i$ be the natural projection for $i = 1, 2$. Then

$$Q_1 := 0 \rightarrow \pi^{-1}(W_1) \rightarrow V \xrightarrow{p_1\pi} W_1 \rightarrow 0$$

and

$$Q_2 := 0 \rightarrow U \rightarrow \pi^{-1}(W_2) \xrightarrow{p_2\pi} W_2 \rightarrow 0$$

are exact sequences and $[[Q]] = [[Q_1]] + [[Q_2]]$ in $G(A)$. $\square$

Furthermore we have $([V], [[Q]]) \geq ([V], [[Q_1]])$ for any module V as $([V], [[E]]) \geq 0$ for any exact sequence E by 5.6(1).

We can prove the next result.

Theorem 5.8. *Let M be an indecomposable R -module and C an indecomposable Γ -module, such that M is a direct summand of C^R with multiplicity n . Then $[[A(M)_\Gamma]] = n \sum_{g \in T(C)} [[A(C_g)]]$.*

Proof. We first show that $([V], [[A(M)_\Gamma]]) - n \sum_{g \in T(C)} ([V], [[A(C_g)]]) = 0$ for all indecomposable Γ -modules V . Using Frobenius reciprocity 5.5, we have $([V], [[A(M)_\Gamma]]) = ([V^R], [[A(M)]])$ which is equal to the multiplicity of M as a direct summand in V^R . By Lemma 5.3 we have that $(M_\Gamma)(C^R)_\Gamma = \bigoplus_{g \in G} C_g$. Therefore M is a direct summand of V^R if and only if $V \cong C_g$ for some $g \in G$. But in this case $V^R = C^R$ and therefore the multiplicity of M as a direct summand of V^R is n . It remains to show that $[[A(M)_\Gamma]]$ is a linear combination of Auslander-Reiten sequences X_i . We have $M_\Gamma = q \sum_{g \in T(C)} C_g$. By 5.7 we know that $[[A(M)_\Gamma]]$ can be written as sum of $[[Q_g^i]]$ where the Q_g^i are exact sequences starting in C_g for $g \in G$ for $1 \leq i \leq q$. Suppose one of them is non-split and not an Auslander-Reiten sequence X_i . By the second part of 5.6 there exists an indecomposable direct summand L of the middle term of some $A(C_g)$ such that $([L], [[A(M)_\Gamma]]) \geq 1$. There is no irreducible map from C_l to C_h for any $l, h \in G$ as both modules have the same dimension. Therefore $L \not\cong C_l$ for all $l \in G$. By 5.6 we have $([L], [[A(C_l)]]) = 0$ for all $l \in G$, which is a contradiction to the first part. $\square$

Clearly if $A(M)$ is an Auslander-Reiten sequence and M a non-projective indecomposable module, then $\tau(M)$, M and the middle term of $A(M)$ have no direct summand in common. The same is true for the restriction of the Auslander-Reiten sequence to Γ .

Lemma 5.9. *Let M be an indecomposable R -module and $A(M) := 0 \rightarrow \tau(M) \rightarrow X \rightarrow M \rightarrow 0$. Then the pair $\tau(M)_\Gamma$ and X_Γ and the pair M_Γ and X_Γ have no direct summand in common. If $M_g \not\cong \tau(M)$, then $\tau(M)_\Gamma$ and M_Γ have no direct summand in common.*

Proof. Suppose there is an indecomposable Γ -module Q , such that $Q|(M_\Gamma)$ and $Q|(X_\Gamma)$. By 5.3 there exists an indecomposable direct summand E of X and a $g \in G$ such that $E \cong M_g$. As M and M_g have the same dimension there is no irreducible map from E to M which is a contradiction. By an analogous argument it is clear that $\tau(M)_\Gamma$ and X_Γ have no direct summand in common. Suppose now that M_Γ and $\tau(M)_\Gamma$ have a common direct summand. Then there exists a $g \in G$ such that $\tau(M) \cong M_g$, which is a contradiction. $\square$

This gives the next

Corollary 5.10. *Let M be an indecomposable, non-projective R -module. Let C be an indecomposable direct summand of M_Γ with multiplicity n . Then M is a direct summand of C^R with multiplicity n . Furthermore if N is the middle term of $A(M)$ and Q the middle term of $A(C)$, then $N_\Gamma = n \bigoplus_{g \in T(C)} Q_g$ and $\tau(M)_\Gamma = n \bigoplus_{g \in T(C)} \tau(C)_g$.*

Proof. Let $A(M) := 0 \rightarrow \tau(M) \rightarrow \bigoplus_{1 \leq i \leq t} d_i N_i \rightarrow M \rightarrow 0$ and $A(C) := 0 \rightarrow \tau(C) \rightarrow \bigoplus_{1 \leq i \leq s} f_i Q_i \rightarrow C \rightarrow 0$ for Q_i, N_i indecomposable and $d_i, f_i \in \mathbb{N}$. We set $Q := \bigoplus_{1 \leq i \leq s} f_i Q_i$ and $N := \bigoplus_{1 \leq i \leq t} d_i N_i$. Then $d_i N_\Gamma^i = b_i \sum_{g \in T(E^i)} E_g^i$, $\tau(M)_\Gamma = a \sum_{g \in T(L)} L_g$ and $M_\Gamma = q \sum_{g \in T(C)} C_g$ for some indecomposable $L, E^i \in \Gamma\text{-mod}$ and $a, q, b_i \in \mathbb{N}$. Then

$$[[A(M)_\Gamma]] = q \sum_{g \in T(C)} [C_g] + a \sum_{g \in T(L)} [L_g] - \sum_{1 \leq i \leq t} b_i \sum_{g \in T(E^i)} [E_g^i].$$

By 5.8 we also have

$$[[A(M)_\Gamma]] = n \sum_{g \in T(C)} [C_g] + n \sum_{g \in T(C)} [\tau(C)_g] - n \sum_{1 \leq i \leq s} f_i \sum_{g \in T(C)} [Q_g^i].$$

We assume that $M_g \not\cong \tau(M)$. Then by 5.9 the set $\{[C_g], g \in T(C)\} \cup_{1 \leq i \leq t} \{[E_g^i], g \in T(E^i)\} \cup \{[L_g], g \in T(L)\}$ is linearly independent. Similarly the set $\{[C_g], g \in T(C)\} \cup_{1 \leq i \leq s} \{[Q_g^i], g \in T(Q^i)\} \cup \{[\tau(C)_g], g \in T(\tau(C))\}$ is linearly independent. We can see this as follows: if $\tau(C) \cong C_h$ for some $h \in G$, then $\sum_{g \in T(C)} \tau(C)_g = \sum_{g \in T(C)} C_g$ which is a contradiction, as the element $[L]$ would not appear as a summand of $[[A(M)_\Gamma]]$ in the second presentation. Also $Q_h^i \not\cong \tau(C)$ and $Q_h^i \not\cong C$ for some $h \in G$ because there is no irreducible map between elements of the same dimension. We compare now the two presentations and use the linear independency of the indecomposable Γ -modules in $G(\Gamma)$. Then $q = n$, $\tau(M)_\Gamma = n \sum_{g \in T(C)} \tau(C)_g$ and $N_\Gamma = n \sum_{g \in T(C)} Q_g$.

Assume now that $M_g \cong \tau(M)$. Then $\tau(M)_\Gamma = q \sum_{g \in T(C)} C_g$. Therefore

$$[[A(M)_\Gamma]] = 2q \sum_{g \in T(C)} [C_g] - \sum_{1 \leq i \leq t} b_i \sum_{g \in T(E^i)} [E_g^i].$$

By 5.9 we have that the set $\{[C_g], g \in T(C)\} \cup_{1 \leq i \leq t} \{[E_g^i], g \in T(E^i)\}$ is linearly independent in $G(\Gamma)$. We have $\tau(C) = C_h$ for some $h \in G$ by comparing summands in the two presentations of $[[A(M)_\Gamma]]$. Then $\sum_{g \in T(C)} \tau(C)_g = \sum_{g \in T(C)} C_g$. Therefore $q = n$, $\tau(M)_\Gamma = n \sum_{g \in T(C)} \tau(C)_g$ and $N_\Gamma = n \sum_{g \in T(C)} Q_g$. $\square$

We can now investigate the relation between periodic R -modules and periodic Γ -modules.

Lemma 5.11 (periodic modules). *The indecomposable R -module M is periodic if and only if M_Γ contains a periodic direct summand.*

Proof. Let Q be an indecomposable direct summand of M_Γ . Then $M_\Gamma = n \bigoplus_{g \in T(Q)} Q_g$ and $\tau(M)_\Gamma = n \bigoplus_{g \in T(Q)} \tau(Q)_g$ by 5.10. Suppose now that Q is τ -periodic with period m . Then Q_g is τ -periodic with period m and $\tau^m(M)_\Gamma \cong M_\Gamma$ as by 5.10 τ and the restriction to Γ commute. This implies that $M^g \cong \tau^m(M)$. As G has finite order, M is periodic. Similarly, if M is τ -periodic with period m , we have $n \bigoplus_{g \in T(Q)} Q_g = M_\Gamma = \tau^m(M)_\Gamma = n \bigoplus_{g \in T(Q)} \tau^m(Q)_g$. Therefore $\tau^m(Q) \cong Q_g$ for some $g \in G$. As G has finite order, Q is τ -periodic. $\square$

Next, we summarize the properties that we need for the following theorems.

Assumption 5.12. *In the following we assume that Γ and R are Frobenius algebras that satisfy (C').*

The following Lemma shows that this could be slightly weakened.

Lemma 5.13. *The algebra R satisfies (C') if and only if the algebra Γ satisfies (C').*

Proof. The proof is analogous to 5.11. $\square$

We also have

Lemma 5.14. *Suppose $\Gamma \neq S$. Let E be a non-projective Γ -module and N a simple R -module. Then $\underline{\text{Hom}}(S, E) \neq 0$ and $\underline{\text{Hom}}(N, E^R) \neq 0$.*

Proof. We consider the map which maps S to $\text{soc } E$. This map does not factor through Γ . Therefore $\underline{\text{Hom}}(S, E) \neq 0$. We have $\text{Hom}(N, E^R) \cong \text{Hom}(N_\Gamma, E) \cong \text{Hom}(S, E)$ by 5.5. As the restriction to Γ and lifting to R preserves projectivity, we have $\underline{\text{Hom}}(N, E^R) = \underline{\text{Hom}}(S, E) \neq 0$. $\square$

We can now investigate the relation between the Auslander-Reiten components of $T_s(\Gamma)$ and the Auslander-Reiten components of $T_s(R)$. We assume for the next two Theorems that 5.12 is satisfied. We denote by $\text{Obj}(\theta)$ the set of all indecomposable modules in an Auslander-Reiten component θ .

Theorem 5.15. (1) $T_s(\Gamma)$ has a component of tree class $\tilde{D}_n$ if and only if $T_s(R)$ has a component of tree class $\tilde{D}_n$.

(2) Suppose $T_s(R)$ has a component of tree class $\tilde{A}_{1,2}$, then $T_s(\Gamma)$ has also a component of tree class $\tilde{A}_{1,2}$.

Proof. (a) Let $\theta \cong \mathbb{Z}[\tilde{D}_n]$ or $\theta \cong \mathbb{Z}[\tilde{A}_{1,2}]$ be a component of $T_s(R)$. We denote the tree class of θ by $\mathcal{T}$. Then by 2.4 θ contains a simple module M and a projective module P is attached to θ . Let Δ be the component in $T_s(\Gamma)$ containing the simple module $S = M_\Gamma$. Then by 5.8 we have $\text{Obj}(\theta_\Gamma) \subset \cup_{g \in G} \text{Obj}(\Delta_g)$ using induction on the distance of a module in θ to M .

As S is contained in Δ and G acts trivially on S , we have $\Delta_g = \Delta$ for all $g \in G$ and therefore $\text{Obj}(\theta_\Gamma) \subset \text{Obj}(\Delta)$. As P is attached to θ , Γ is attached to Δ . Then Ω induces a fix point free automorphism on the tree class T of Δ .

By 3.3 and 3.11, $\text{mod } R$ has a periodic module that does not lie in θ and therefore by 5.11, we have that $\text{mod } \Gamma$ also contains a periodic module that does not lie in Δ . Using 5.14 we can define a subbaditive, non-zero function on the component Δ as in [ES, 3.2]. The tree class of Δ is therefore in the HPR-list (this is a list of trees given in [HPR, p.286]).

As we have a fix point free automorphism operating on T , this gives $T = A_\infty^\infty$, $T = \tilde{D}_m$ for m odd or $T = \tilde{A}_{1,2}$.

Note that S and Γ/S do not lie in the same τ -orbit, because by 2.1 we would have $2 = l(\Gamma)$. But then S would be periodic which is a contradiction to the fact that M is not periodic.

Twisting with $g \in G$ also induces an automorphism of finite order on Δ , that fixes S and Γ/S . If $T = \tilde{D}_m$, then S and Γ/S have only one predecessor and their predecessors do not lie in the same τ -orbit by 3.5. Therefore g acts as the identity and there are no modules contained in Δ that are twists of each other.

This means that we can embed Δ into θ using induction on the distance of a module in Δ to S . We give a sketch of how to construct this injection: We map S to M . Let $W \in \text{Obj}(\Delta)$ and suppose that there is an arrow from W to S in Δ . Then by 5.10 there exists an R -module J such that J is a summand of the middle term of $A(M)$ and $W|J_\Gamma$. We map W to J . This gives an injection, as for two indecomposable Γ modules W^1 and W^2 with $W^1|Z_\Gamma$ and $W^2|Z_\Gamma$ for an indecomposable R -module Z , we have $W^1 \cong (W^2)_g$ for some $g \in G$. So if W^1 and W^2 lie in Δ we have $W^1 \cong W^2$.

As this embedding respects τ , it induces an embedding $T \subset \mathcal{T}$. Therefore $T = \mathcal{T}$. This proves the first direction of (1) and part (2).

(b) Suppose now that $\Delta \cong \mathbb{Z}[\tilde{D}_n]$, then $S \in \Delta$ and Γ is attached to Δ . Let θ be a component of $T_s(R)$ that contains a simple module M . As in (a) we have $\text{Obj}(\theta_\Gamma) \subset \cup_{g \in G} \text{Obj}(\Delta_g)$.

By 3.11, $\text{mod } \Gamma$ contains a periodic module E that is not in Δ and by 5.11 all direct summands of E^R are periodic and do not lie in θ . As in the first part of the proof, this shows that the tree class T of θ is in the HPR-list.

As $\bar{\alpha}(S) = \bar{\alpha}(\Gamma/S) = 1$ by 3.5 and do not have the same predecessor, g acts as the identity on Δ . That means $\text{Obj}(\theta_\Gamma) \subset \text{Obj}(\Delta)$.

As in (a) we can embed Δ into θ . As τ and the restriction to Γ commute by 5.10, this gives an embedding of $\tilde{D}_n$ into T . Therefore $T = \tilde{D}_n$. $\square$

Note that the embedding of Δ into θ does not respect labels of arrows. The next example shows that part (2) of the previous theorem does not hold in the converse direction.

Example 5.16. Let k be a field of characteristic 2, $\Gamma = kV_4$ and $R = kA_4 \cong kV_4 \rtimes kC_3$. Then Γ has a component with tree class $\tilde{A}_{1,2}$ and R has a component with reduced graph $\tilde{A}_5$ which corresponds to a tree class A_∞ . By 3.3 $T_s(R)$ has no component of Euclidean tree class.

We can also show the following

Theorem 5.17. Suppose $T_s(\Gamma)$ has a component Δ of tree class $\tilde{A}_{1,2}$, then $T_s(R)$ has also a component θ of tree class $T = \tilde{A}_{1,2}$ or $T = A_\infty$. In the second case $\theta \cong \mathbb{Z}[\tilde{A}_n]$, where $n + 1/2$ divides the order of G .

Proof. Let M be a simple R -module and let θ be the component that contains M . As in the proof of the previous Theorem the tree class T of θ is from the HPR-list and θ is not a periodic component. Let Q be the middle term of $A(M)$. Then $Q_\Gamma = N \oplus N$ for some indecomposable Γ -module N .

Suppose first that $Q \cong L \oplus L_g$ for L an indecomposable R -module and $g \in G$ such that $L_\Gamma \cong N$ and $L_g \not\cong L$. Then $\bar{\alpha}(M) = 2$ and g acts as a graph automorphism on T of finite order and is not the identity. Also this automorphism commutes with τ . Furthermore the middle term of $A(\tau^{-1}(L))$ is $M \oplus M_{g^{-1}}$. As $M \not\cong M_{g^{-1}}$ we have $\theta \cong \mathbb{Z}[\tilde{A}_n]$, where $n = 2|g| - 1$.

Suppose now that Q is indecomposable. Then either $M \oplus M$ or $M \oplus M_g$ is the non-projective summand in the middle term of $A(\tau^{-1}(Q))$. In the first case, we have $\bar{\alpha}(M) = \bar{\alpha}(Q) = 1$. Therefore $T = Q \xrightarrow{(1,2)} M$. This contradicts with the HPR-list. In the second case we have $\bar{\alpha}(Q) = 2$ and $\bar{\alpha}(M) = \bar{\alpha}(M_g) = 1$. As all three are in different τ -orbits, we have $T = M \longrightarrow Q \longleftarrow M_g$ which is also a contradiction, as Δ is not of finite type.

Suppose now that $Q \cong L \oplus L$. Then either $M \oplus M$ or $M \oplus M_g$ is the non-projective summand in the middle term of $A(\tau^{-1}(L))$. The first case

gives $\theta \cong \mathbb{Z}[A_{1,2}]$. In the second case we have $T = \tilde{B}_2$, which we can exclude using 3.3. $\square$

REFERENCES

- [ASS] I. Assem, D. Simson and A. Skowroński, *Elements of the Representation Theory of Associative Algebras 1: Techniques of Representation Theory*, London Mathematical Society Student Texts, 65. Cambridge University Press, Cambridge, 2006.
- [Ben1] D. Benson, *Representations and Cohomology 1: Basic Representation Theory of Finite Groups and Associative Algebras*, Cambridge Studies in Advanced Mathematics, 30. Cambridge University Press, Cambridge, 1998.
- [BP] D. Benson and R. Parker, *The Green ring of a finite group*, J. Algebra 87 (1984), no. 2, 290-331.
- [BR] M. C. R. Butler and C. M. Ringel, *Auslander-Reiten sequences with few middle terms and applications to string algebras*, Comm. Algebra 15 (1987), no. 1-2, 145-179.
- [E] K. Erdmann, *The Auslander-Reiten quiver of restricted enveloping algebras*. Representation Theory of Algebras, CMS Conf. Proc. 18, Amer. Math. Soc., Providence, RI, 1996, 201-214.
- [ES] K. Erdmann and A. Skowroński, *On Auslander-Reiten components of blocks and self-injective biserial algebras*, Trans. Amer. Math. Soc. 330 (1992), no. 1, 165-189.
- [F1] R. Farnsteiner, *Periodicity and representation type of modular Lie algebras*, J. Reine Angew. Math. 464 (1995), 47-65.
- [F2] R. Farnsteiner, *On Auslander-Reiten quivers of enveloping algebras of restricted Lie algebras*, Math. Nachr. 202 (1999), 43-66.
- [F3] R. Farnsteiner, *Representations of blocks associated to induced modules of restricted Lie algebras*, Math. Nachr. 179 (1996), 57-88.
- [FS] R. Farnsteiner and H. Strade, *Modular Lie Algebras and Their Representations*, Monographs and Textbooks in Pure and Applied Mathematics, 116. Marcel Dekker, Inc., New York, 1988.
- [FS1] R. Farnsteiner and H. Strade, *Shapiro's lemma and its consequences in the cohomology theory of modular Lie algebras*, Math. Z. 206 (1991), no. 1, 153-168.
- [HPR] D. Happel, U. Preiser and C. M. Ringel, *Vinberg's characterization of Dynkin diagrams using subadditive functions with application to DTr-periodic modules*. Representation Theory II, Lecture Notes in Math. 832 (1980), Springer, Berlin, 1980, 280-294.
- [SSS] M. Schaps, D. Shapira and O. Shlomo, *Quivers of blocks with normal defect group*, Proc. Sympos. Pure Math. 63, Amer. Math. Soc. Providence, RI, 1998.
- [SM] S. O. Smalø, *Local limitations of the Ext functor do not exist*, Bull. London Math. Soc. 38 (2006), 97-98.
- [Sk] A. Skowroński, *Selfinjective algebras: finite and tame type*. Trends in Representation Theory of Algebras and Related Topics. Contemporary Math. 406 (2006), Amer. Math. Soc. Providence, RI, 169-238.
- [W] P. Webb, *The Auslander-Reiten quiver of a finite group*, Math. Z. 179 (1982), no. 1, 97-121.

SARAH SCHEROTZKE

MATHEMATICAL INSTITUTE, UNIVERSITY OF OXFORD, 24-29 ST. GILES, OXFORD OX1 3LB, UNITED KINGDOM

E-mail address: `scherotz@maths.ox.ac.uk`